## Asthma Treatment Plan

## CITY OF AUGUSTA CHILDCARE BUREAU

(Please Print)				Date of Birth Effective Date			
ivame							
Doctor			Parent/Guardian (if applicable) Emergency Contact		ency Contact		
Phone			Phone		Phone	Phone	
HEALTH	Y		Take daily medicin to be used with spa		d dose	inhalers (MDI)	
		ve <u>all</u> of these: fing is good	MEDICINE HOW MUCH to take and HOW OFTEN to take it			Triggers Check all items	
	<ul> <li>No cough or wheeze</li> <li>Sleep through the night</li> <li>Can work, exercise, and play</li> </ul>		☐ Advair® 100, 250, 500				that trigger patient's asthma:  Chalk dust Cigarette Smoke Second hand Smoke Colds/Flu Dust mites, dust, stuffed animals, carpet Exercise Mold
And/or Peak flow above			Remember to rinse your mouth after taking inhaled medicine.				Ozone alert days
lf ex	vercise tri	ngers vour asthma	Remember a, take this medicine	to rinse your mouth a		ng inhaled medicine. utes before exercise.	<ul><li>Pests - rodents cockroaches</li></ul>
			Continue daily medicine(s) and add fast-acting medicine(s).				<ul><li>□ Pets - animal dander</li><li>□ Plants, flowers,</li></ul>
		ve <u>any</u> of these:					cut grass, polle Strong odors,
	Exposure to known trigger     Cough     Mild wheeze     Tight chest     Coughing at night     Other:		☐ Accuneb® 0.63, 1.25 mg 1 unit nebulized every 4 hours as needed ☐ Albuterol 1.25, 2.5 mg 1 unit nebulized every 4 hours as needed ☐ Albuterol ☐ Pro-Air ☐ Proventil® .2 puffs MDI every 4 hours as needed ☐ Ventolin® ☐ Maxair ☐ Xopenex® .2 puffs MDI every 4 hours as needed ☐ Xopenex® 0.31, 0.63, 1.25 mg 1 unit nebulized every 4 hours as needed ☐ Increase the dose of, or add:				perfumes, clearing products, scented products, scented products and the summer of the
And/or Peak flo	ow from	to to	If fast-acting medicine is needed more than 2 times a week, except before exercise, then call your doctor.			Other:	
EMERG	Your ast getting • Fast-act	thma is worse fast: ing medicine did not	Take these medicines NOW and call 911.  Asthma can be a life-threatening illness. Do not wait!  Accuneb® 0.63, 1.25 mg 1 unit nebulized every 20 minutes  Albuterol 1.25, 2.5 mg 1 unit nebulized every 20 minutes  Albuterol Pro-Air Proventil® .2 puffs MDI every 20 minutes  Ventolin® Maxair Xopenex® 2 puffs MDI every 20 minutes  Xopenex® 0.31, 0.63, 1.25 mg 1 unit nebulized every 20 minutes  Other			and other.	
And/or Peak	<ul><li>Breathin</li><li>Nose op</li><li>Ribs sho</li><li>Trouble</li><li>Lips blu</li></ul>	ow walking and talking e • Fingernails blue				This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.	
		the proper method medications named	able and has been instructed in of self-administering of the inhaled	PARENT/GUARDIAN SIGN	IATURE	ugedezes (gazt angeleg sich zu entschap gewossen end gefall die Stadt zu Schmiederschaft zu der Schmieder	